

## Tutorial\*

### Tutorial 25: Calculating Mixtures of Local Anesthetic and Morphine for Implantable Intrathecal Pumps

Glen Atlas, M.D., M.Sc.

Department of Anesthesiology, Sebastian River Medical Center, 13695 U.S. Highway 1, Sebastian, Florida 32958, U.S.A.

The use of implantable intrathecal pumps has become increasingly more complex with clinicians' combining opiates and local anesthetics such as morphine and bupivacaine. This technique produces heightened analgesia and is especially useful for the treatment of cancer-related neuropathic pain [1].

In this patient population, use of low-dose intrathecal bupivacaine does not appear to cause significant neurologic sequelae [2]. However, use of spinal catheters, with high concentrations of lidocaine, has been associated with cauda equina syndrome [3]. Recently, successful long-term use of intracisternal bupivacaine has been described for chronic facial pain [4,5]. It is conceivable that the combination of intrathecal bupivacaine and morphine might be useful in treating recalcitrant chronic regional pain syndromes such as reflex sympathetic dystrophy or causalgia.

Potential side effects of intrathecal local anesthetics including hypotension, bradycardia, and motor weakness should be evaluated [6]. In addition, the side effects of intrathecal opiates, such as respiratory depression, pruritic, and urinary retention must also be recognized [7]. If necessary, the reservoir and intrathecal catheter of these pumps can be easily drained.

The following sets of equations allow for readily available concentrations of both morphine and bupivacaine to be mixed so that a desired total daily dose of each can be administered. Two sets of equations will be described: one for pumps which are externally programmable, such as the Medtronic, and another for fixed-rate pumps, such as the Infusaid or Arrow. Table 1 provides a listing of terms and their definitions, along with appropriate units of measure.

#### Determining Mixtures for Programmable Intrathecal Pumps

Initially, a dimensionless constant R is defined which equals the ratio of the total daily dose of local anesthetic,  $T_{LA}$ , to the

total daily dose of morphine,  $T_{MSO_4}$ . This constant is also equivalent to the ratio of the final concentration of local anesthetic,  $C_{LA-final}$ , to the final concentration of morphine,  $C_{MSO_4-final}$ .

$$R = \frac{C_{LA-final}}{C_{MSO_4-final}} = \frac{\text{Total daily dose of local anesthetic in mg/day}}{\text{Total daily dose of morphine in mg/day}} = \frac{T_{LA}}{T_{MSO_4}} \quad (1)$$

The final concentration of morphine,  $C_{MSO_4-final}$ , can then be determined using the initial concentration of local anesthetic,  $C_{LA-initial}$ , and the initial concentration of morphine,  $C_{MSO_4-initial}$ .

$$C_{MSO_4-final} = \frac{1}{\frac{R}{C_{LA-initial}} + \frac{1}{C_{MSO_4-initial}}} \quad (2)$$

By using R and  $C_{MSO_4-final}$  from equation 2, the final concentration of local anesthetic,  $C_{LA-final}$ , can be found:

$$C_{LA-final} = R \cdot C_{MSO_4-final} \quad (3)$$

The volumes of morphine,  $\text{Volume}_{MSO_4-initial}$ , and local anesthetic,  $\text{Volume}_{LA-initial}$ , to be drawn up, mixed, and infused into the pump reservoir are then calculated [8]:

$$\text{Volume}_{MSO_4-initial} = \frac{C_{MSO_4-final} \cdot \text{Volume}_{reservoir}}{C_{MSO_4-initial}} \quad (4)$$

and

$$\text{Volume}_{LA-initial} = \frac{C_{LA-final} \cdot \text{Volume}_{reservoir}}{C_{LA-initial}} \quad (5)$$

The pump flow rate can then be determined:

$$\text{Pump flow rate in ml/day} = \frac{T_{MSO_4}}{C_{MSO_4-final}} \quad (6)$$

\*The purpose of this section is to provide the reader with a series of tutorial lessons that may be used for self-study in pain medicine. CME credits will not be granted for the completion of this tutorial.

†See "Formula Derivations" section. Volumes and concentrations are expressed in milliliter and milligram per milliliter, respectively.

Table 1. List of terms.

Term	Meaning	Units
$C_{LA-initial}$	Initial concentration of local anesthetic	mg/ml
$C_{MSO_4-initial}$	Initial concentration of morphine	mg/ml
$C_{LA-final}$	Final concentration of local anesthetic	mg/ml
$C_{MSO_4-final}$	Final concentration of morphine	mg/ml
$Volume_{LA-initial}$	Initial volume of local anesthetic	ml
$Volume_{MSO_4-initial}$	Initial volume of morphine	ml
$Volume_{LA-final}$	Final volume of local anesthetic	ml
$Volume_{MSO_4-final}$	Final volume of morphine	ml
$T_{LA}$	Total daily dose of local anesthetic	mg/day
$T_{MSO_4}$	Total daily dose of morphine	mg/day
$F_p$	Flow rate for fixed-rate pumps	ml/day
$R$	$\frac{C_{LA-final}}{C_{MSO_4-final}}$ or $\frac{T_{LA}}{T_{MSO_4}}$	dimensionless
$Volume_{NS}$	Volume of preservative-free normal saline	ml

or

$$\text{Pump flow rate in ml/day} = \frac{T_{LA}}{C_{LA-final}} \quad (7)$$

Calculating both Equations 6 and 7 is useful in checking for accuracy.

The duration that the pump reservoir can be used, until it requires refilling, can be easily found:

$$\text{Duration in days} = \frac{\text{Volume}_{\text{reservoir}}}{\text{Pump flow rate}} \quad (8)$$

### Determining Mixtures for Fixed-Rate Intrathecal Pumps

For fixed-rate pumps, such as the Infusaid, the final concentrations of morphine and local anesthetic can be readily determined:

$$C_{MSO_4-final} = \frac{T_{MSO_4}}{F_p} \quad (9)$$

and

$$C_{LA-final} = \frac{T_{LA}}{F_p} \quad (10)$$

$F_p$  represents the fixed flow rate for the pump.

The initial volumes of local anesthetic and morphine to be drawn up and mixed can then be determined (see "Formula Derivations"):

$$\text{Volume}_{LA-initial} = \frac{T_{LA}}{F_p} \cdot \frac{\text{Volume}_{\text{reservoir}}}{C_{LA-initial}} \quad (11)$$

and

$$\text{Volume}_{MSO_4-initial} = \frac{T_{MSO_4}}{F_p} \cdot \frac{\text{Volume}_{\text{reservoir}}}{C_{MSO_4-initial}} \quad (12)$$

Preservative-free normal saline is then added to the above volumes of local anesthetic and morphine so that the final total volume is equal to that of the reservoir. This volume can be calculated:

$$\text{Volume}_{NS} = \text{Volume}_{\text{reservoir}} - (\text{Volume}_{LA-initial} + \text{Volume}_{MSO_4-initial}) \quad (13)$$

Obviously, if  $Volume_{NS}$  is less than zero, then either  $C_{LA-initial}$  or  $C_{MSO_4-initial}$  or both must be increased to reduce  $Volume_{LA-initial}$  or  $Volume_{MSO_4-initial}$ , respectively.

### Checking Accuracy

The above calculations, for both programmable and fixed-rate pumps, can be checked by multiplying the final concentrations, of the local anesthetic and morphine, by the pump flow rate.

$$C_{LA-final} \cdot (\text{Pump flow rate}) = T_{LA} \quad (14)$$

and

$$C_{MSO_4-final} \cdot (\text{Pump flow rate}) = T_{MSO_4} \quad (15)$$

The following equations are also necessary in accuracy checking:

$$C_{LA-initial} \cdot \text{Volume}_{LA-initial} = C_{LA-final} \cdot \text{Volume}_{LA-final} \quad (16)$$

and

$$C_{MSO_4-initial} \cdot \text{Volume}_{MSO_4-initial} = C_{MSO_4-final} \cdot \text{Volume}_{MSO_4-final} \quad (17)$$

### Examples

Two calculations using the preceding formulas follow.

#### Example 1

It is desired to deliver 4 mg/day of intrathecal morphine with 5 mg/day of intrathecal bupivacaine via a *programmable* pump. The initial concentration of morphine,  $C_{MSO_4-initial}$ , is 25.0 mg/ml; and the initial concentration of bupivacaine,  $C_{LA-initial}$ , is 7.5 mg/ml. The pump reservoir has a capacity of 18 ml. Using Equation 1:

$$R = \frac{5}{4}$$

The final concentration of morphine,  $C_{MSO_4-final}$ , can be obtained from Equation 2:

$$C_{MSO_4-final} = \frac{1}{\left(\frac{5}{4}\right) + \frac{1}{25}} = 4.84 \text{ mg/ml}$$

Using Equation 3, the final concentration of bupivacaine can be found:

$$C_{LA-final} = R \cdot C_{MSO_4-final} = \frac{5}{4} \cdot (4.84) = 6.05 \text{ mg/ml}$$

The volumes of morphine and local anesthetic, to be drawn up and mixed, are determined from Equations 4 and 5:

$$\text{Volume}_{MSO_4-initial} = \frac{(4.84) \cdot 18}{25} = 3.48 \text{ ml}$$

$$\text{Volume}_{LA-initial} = \frac{(6.05) \cdot 18}{7.5} = 14.52 \text{ ml}$$

The pump flow rate is then calculated from Equations 6 or 7:

$$\text{Pump flow rate} = \frac{4}{4.84} = \frac{5}{6.05} = 0.826 \text{ ml/day}$$

From Equation 8, the duration that the pump can be used until its reservoir needs to be refilled is:

$$\text{Duration in days} = \frac{18}{0.826} = 21.8 \text{ days}$$

The accuracy can be checked by using Equations 14 and 15 to verify the total daily dose of local anesthetic and morphine.

$$C_{\text{LA-final}} \cdot (\text{pump flow rate}) = 6.05 \cdot 0.826 = 5 \text{ mg/day}$$

and

$$C_{\text{MSO}_4\text{-final}} \cdot (\text{pump flow rate}) = 4.84 \cdot 0.826 = 4 \text{ mg/day}$$

The initial and final concentrations and volumes can also be checked by using Equations 16 and 17.

$$C_{\text{LA-initial}} \cdot \text{Volume}_{\text{LA-initial}} = 7.5 \cdot 14.52 = 109 \text{ mg}$$

$$C_{\text{LA-final}} \cdot \text{Volume}_{\text{LA-final}} = 6.05 \cdot 18 = 109 \text{ mg}$$

and

$$C_{\text{MSO}_4\text{-initial}} \cdot \text{Volume}_{\text{MSO}_4\text{-initial}} = 25 \cdot 3.48 = 87 \text{ mg}$$

$$C_{\text{MSO}_4\text{-final}} \cdot \text{Volume}_{\text{MSO}_4\text{-final}} = 4.84 \cdot 18 = 87 \text{ mg}$$

### Example 2

It is desired to deliver 7 mg/day of intrathecal morphine and 5 mg/day of intrathecal bupivacaine via a *fixed-rate* pump with a reservoir volume of 50 ml. The pump has a preset flow rate of 2 ml/day. The initial concentration of local anesthetic,  $C_{\text{LA-initial}}$ , is 7.5 mg/ml; and the initial concentration of morphine,  $C_{\text{MSO}_4\text{-initial}}$ , is 10.0 mg/ml.

The final concentrations of morphine and local anesthetic can be determined from Equations 9 and 10.

$$C_{\text{MSO}_4\text{-final}} = \frac{7}{2} = 3.5 \text{ mg/ml}$$

and

$$C_{\text{LA-final}} = \frac{5}{2} = 2.5 \text{ mg/ml}$$

The initial volumes of local anesthetic and morphine to be drawn up and mixed can be determined using Equations 11 and 12.

$$\text{Volume}_{\text{LA-initial}} = \frac{5}{2} \cdot \frac{50}{7.5} = 16.67 \text{ ml}$$

and

$$\text{Volume}_{\text{MSO}_4\text{-initial}} = \frac{7}{2} \cdot \frac{50}{10} = 17.5 \text{ ml}$$

The volume of preservative-free normal saline to be drawn up and mixed to the above volumes is determined from Equation 13.

$$\text{Volume}_{\text{NS}} = 50 - (16.67 + 17.5) = 15.83 \text{ ml}$$

The accuracy can be checked by using Equations 14 and 15 to verify the total daily dose of local anesthetic and morphine.

$$C_{\text{LA-final}} \cdot (\text{pump flow rate}) = 2.5 \cdot 2.0 = 5 \text{ mg/day}$$

and

$$C_{\text{MSO}_4\text{-final}} \cdot (\text{pump flow rate}) = 3.5 \cdot 2.0 = 7 \text{ mg/day}$$

The initial and final concentrations and volumes can also be checked by using Equations 16 and 17.

$$C_{\text{LA-initial}} \cdot \text{Volume}_{\text{LA-initial}} = 7.5 \cdot 16.67 = 125 \text{ mg}$$

$$C_{\text{LA-final}} \cdot \text{Volume}_{\text{LA-final}} = 2.5 \cdot 50 = 125 \text{ mg}$$

and

$$C_{\text{MSO}_4\text{-initial}} \cdot \text{Volume}_{\text{MSO}_4\text{-initial}} = 10 \cdot 17.5 = 175 \text{ mg}$$

$$C_{\text{MSO}_4\text{-final}} \cdot \text{Volume}_{\text{MSO}_4\text{-final}} = 3.5 \cdot 50 = 175 \text{ mg}$$

### Formula Derivations

Equation 2 is derived from the following formulas:

$$C_{\text{MSO}_4\text{-initial}} \cdot \text{Volume}_{\text{MSO}_4\text{-initial}} = C_{\text{MSO}_4\text{-final}} \cdot \text{Volume}_{\text{MSO}_4\text{-final}} \quad (1)$$

and

$$C_{\text{LA-initial}} \cdot \text{Volume}_{\text{LA-initial}} = C_{\text{LA-final}} \cdot \text{Volume}_{\text{LA-final}} \quad (2)$$

Realizing that:

$$\text{Volume}_{\text{MSO}_4\text{-initial}} + \text{Volume}_{\text{LA-initial}} = \text{Volume}_{\text{reservoir}} \quad (3)$$

and

$$\text{Volume}_{\text{MSO}_4\text{-final}} = \text{Volume}_{\text{LA-final}} = \text{Volume}_{\text{reservoir}} \quad (4)$$

Equations 1 and 2 can then be rearranged, combined as in (3), and expressed using (4) as:

$$\frac{C_{\text{MSO}_4\text{-final}} \cdot \text{Volume}_{\text{reservoir}}}{C_{\text{MSO}_4\text{-initial}}} + \frac{C_{\text{LA-final}} \cdot \text{Volume}_{\text{reservoir}}}{C_{\text{LA-initial}}} = \text{Volume}_{\text{reservoir}} \quad (5)$$

Note that  $\text{Volume}_{\text{reservoir}}$  can then be canceled from each of the above terms. In addition,  $C_{\text{LA-final}}$  can be substituted, with Equation 3, and a common denominator obtained.

$$\frac{C_{\text{MSO}_4\text{-final}} \cdot C_{\text{LA-initial}} + R \cdot C_{\text{MSO}_4\text{-final}} \cdot C_{\text{MSO}_4\text{-initial}}}{C_{\text{MSO}_4\text{-initial}} \cdot C_{\text{LA-initial}}} = 1 \quad (6)$$

Solving for  $C_{\text{MSO}_4\text{-final}}$  yields Equation 2:

$$C_{\text{MSO}_4\text{-final}} = \frac{1}{\left( \frac{R}{C_{\text{LA-initial}}} + \frac{1}{C_{\text{MSO}_4\text{-initial}}} \right)} \quad (7)$$

Equations 11 and 12 are derived from Equations 1 and 2, which are rearranged:

$$\text{Volume}_{\text{MSO}_4\text{-initial}} = \frac{C_{\text{MSO}_4\text{-final}} \cdot \text{Volume}_{\text{reservoir}}}{C_{\text{MSO}_4\text{-initial}}} \quad (8)$$

and

$$\text{Volume}_{\text{LA-initial}} = \frac{C_{\text{LA-final}} \cdot \text{Volume}_{\text{reservoir}}}{C_{\text{LA-initial}}} \quad (9)$$

Substituting Equation 9,  $C_{\text{MSO}_4\text{-final}} = \frac{T_{\text{MSO}_4}}{F_p}$  and Equation 10,  $C_{\text{LA-final}} = \frac{T_{\text{LA}}}{F_p}$  yields Equations 11 and 12:

**Table 2.** Software program, written in Microsoft QBasic, executes the described equations for programmable intrathecal pumps.

```

CLS
format$ = "##.###"
INPUT "Enter the total daily dose of local anesthetic in mg"; TDLA
INPUT "Enter the total daily dose of morphine in mg"; TDMS
INPUT "Enter the initial concentration of local anesthetic in mg/ml"; CLAI
INPUT "Enter the initial concentration of morphine in mg/ml"; CMSI
INPUT "Enter the volume of the pump reservoir in ml"; VolRes
,
R = TDLA / TDMS
CMSF = 1 / ((R / CLAI) + (1 / CMSI))
CLAF = R * CMSF
VolMS = (CMSF * VolRes) / CMSI
VolLA = (CLAF * VolRes) / CLAI
PumpflowrateLA = TDLA / CLAF
PumpflowrateMS = TDMS / CMSF
,
LPRINT "Enter the total daily dose of local anesthetic in mg: "; USING format$; TDLA
LPRINT "Enter the total daily dose of morphine in mg: "; USING format$; TDMS
LPRINT "Enter the initial concentration of local anesthetic in mg/ml: "; USING format$; CLAI
LPRINT "Enter the initial concentration of morphine in mg/ml: "; USING format$; CMSI
LPRINT "Enter the volume of the pump reservoir in ml: "; USING format$; VolRes
LPRINT
LPRINT
LPRINT "Volume of morphine to be drawn up in ml: "; USING format$; VolMS
LPRINT "Volume of local anesthetic to be drawn up in ml: "; USING format$; VolLA
LPRINT "Final concentration of morphine to be infused in mg/ml: "; USING format$; CMSF
LPRINT "Final concentration of local anesthetic to be infused in mg/ml: "; USING format$; CLAF
LPRINT "Pump flow rate in ml/day: "; USING format$; PumpflowrateLA
LPRINT "Pump flow rate in ml/day (check): "; USING format$; PumpflowrateMS
LPRINT "Duration of mixture in reservoir in days: "; USING format$; (VolRes / PumpflowrateLA)
LPRINT
LPRINT
LPRINT "Check: Total daily dose of local anesthetic = CLAF * Pump flow rate ="; CLAF * PumpflowrateLA; "mg"
LPRINT "Check: Total daily dose of morphine = CMSF * Pump flow rate ="; CMSF * PumpflowrateMS; "mg"
LPRINT "Check: CLAI*VolLA ="; CLAI * VolLA; "mg "; "CLAF*VolRes ="; CLAF * VolRes; "mg"
LPRINT "Check: CMSI*VolMS ="; CMSI * VolMS; "mg "; "CMSF*VolRes ="; CMSF * VolRes; "mg"
LPRINT CHR$(12)
SYSTEM
END

```

**Table 3.** Software program, written in Microsoft QBasic, executes the described equations for fixed-rate intrathecal pumps.

```

CLS
format$ = "##.###"
INPUT "Enter the total daily dose of local anesthetic in mg"; TDLA
INPUT "Enter the total daily dose of morphine in mg"; TDMS
INPUT "Enter the initial concentration of local anesthetic in mg/ml"; CLAI
INPUT "Enter the initial concentration of morphine in mg/ml"; CMSI
INPUT "Enter the volume of the pump reservoir in ml"; VolRes
INPUT "Enter the flow rate of the pump in ml/day"; FRP
,
CMSF = TDMS / FRP
CLAF = TDLA / FRP
VolMS = (CMSF * VolRes) / CMSI
VolLA = (CLAF * VolRes) / CLAI
VolNS = VolRes - (VolMS + VolLA)
,
LPRINT "Enter the total daily dose of local anesthetic in mg: "; USING format$; TDLA
LPRINT "Enter the total daily dose of morphine in mg: "; USING format$; TDMS
LPRINT "Enter the initial concentration of local anesthetic in mg/ml: "; USING format$; CLAI
LPRINT "Enter the initial concentration of morphine in mg/ml: "; USING format$; CMSI

```

(Table continues on next page)

**Table 3.** Software program, written in Microsoft QBasic, executes the described equations for fixed-rate intrathecal pumps. (Continued)

```

LPRINT "Enter the volume of the pump reservoir in ml: "; USING format$; VolRes
LPRINT "Enter the flow rate of the pump in ml/day: "; USING format$; FRP
LPRINT
LPRINT
LPRINT "Volume of morphine to be drawn up in ml: "; USING format$; VolMS
LPRINT "Volume of local anesthetic to be drawn up in ml: "; USING format$; VolLA
LPRINT "Volume of normal saline to be drawn up in ml: "; USING format$; VolNS
LPRINT "Final concentration of morphine to be infused in mg/ml: "; USING format$; CMSF
LPRINT "Final concentration of local anesthetic to be infused in mg/ml: "; USING format$; CLAF
LPRINT "Duration of mixture in reservoir in days: "; USING format$; (VolRes / FRP)
LPRINT
LPRINT
LPRINT "Check: Total daily dose of local anesthetic = CLAF * pump flow rate ="; CLAF * FRP; "mg"
LPRINT "Check: Total daily dose of morphine = CMSF * pump flow rate ="; CMSF * FRP; "mg"
LPRINT "Check: CLAI*VolLA ="; CLAI * VolLA; "mg "; "CLAF*VolRes ="; CLAF * VolRes; "mg"
LPRINT "Check: CMSI*VolMS ="; CMSI * VolMS; "mg "; "CMSF*VolRes ="; CMSF * VolRes; "mg"
LPRINT CHR$(12)
SYSTEM
END

```

$$\text{Volume}_{\text{LA-initial}} = \frac{T_{\text{LA}}}{F_p} \cdot \frac{\text{Volume}_{\text{reservoir}}}{C_{\text{LA-initial}}} \quad (10)$$

and

$$\text{Volume}_{\text{MSO}_4\text{-initial}} = \frac{T_{\text{MSO}_4}}{F_p} \cdot \frac{\text{Volume}_{\text{reservoir}}}{C_{\text{MSO}_4\text{-initial}}} \quad (11)$$

### Summary

The clinician will find the preceding sets of equations useful in the day-to-day management of morphine and local anesthetic combinations for administration via implantable intrathecal pumps. Other opiate and local anesthetic combinations, with the appropriate changes in concentrations and nomenclature, can be substituted. In addition, the software programs in Tables 2 and 3, written in Microsoft QBasic, allow for the rapid computation of the necessary final parameters.

### References

1. Krames E: The chronic intraspinal use of opiate and local anesthetic mixtures for the relief of intractable pain: When all else fails! *Pain* 1993;55:1-4
2. Sjöberg M, Karlsson PA, Nordberg C: Neuropathologic findings after long-term intrathecal infusion of morphine and bupivacaine for pain treatment in cancer patients. *Anesthesiology* 1992;76:173-186
3. Rigler ML, Drasner K, Krejcie TC et al: Cauda equina syndrome after continuous spinal anesthesia. *Anesthesia and Analgesia* 1995;72:275-281
4. Appलगren L, Janson M, Nitescu P et al: Continuous intracisternal and high cervical intrathecal bupivacaine analgesia in refractory head and neck pain. *Anesthesiology* 1996;84:256-272
5. Carpenter R, Rauck R: Refractory head and neck pain: A difficult problem and a new alternative therapy. *Anesthesiology* 1996;84:249-252
6. Bridenbaugh PO: Complications of local anesthetic neural blockade. In Bridenbaugh PO, Cousins MJ (eds): *Neural Blockade in Clinical Anesthesia and Pain Management* (2nd ed). Philadelphia, JB Lippincott, 1988, pp 695-735
7. Waldman SD, Leak DW, Kennedy LD et al: Intraspinal opioid therapy. In Patt RB (ed): *Cancer Pain*. Philadelphia, JB Lippincott, 1993, pp 285-328
8. Atlas G: A method to quickly calculate mixtures for epidural infusions. *Anesthesia and Analgesia* 1995;80:642

## Appendix: Continuing Medical Education Questions

For a *programmable* intrathecal pump, it is desired to deliver 3 mg per day of bupivacaine and 5 mg per day of morphine. The pump reservoir has a capacity of 18 ml. The initial concentration of local anesthetic,  $C_{LA-initial}$ , is 7.5 mg/ml and the initial concentration of morphine,  $C_{MSO_4-initial}$ , is 25.0 mg/ml.

- Using Equation 1, determine R.
  - 0.4
  - 0.5
  - 0.6
  - 0.7
- Find the final concentration of morphine,  $C_{MSO_4-final}$ , from Equation 2.
  - 7.33 mg/ml
  - 8.33 mg/ml
  - 9.33 mg/ml
  - 10.33 mg/ml
- Calculate the final concentration of local anesthetic,  $C_{LA-final}$ , from Equation 3.
  - 2 mg/ml
  - 3 mg/ml
  - 4 mg/ml
  - 5 mg/ml
- Using Equation 4, find the initial volume of morphine,  $Volume_{MSO_4-initial}$ .
  - 6.0 ml
  - 7.0 ml
  - 8.0 ml
  - 9.0 ml
- Using Equation 5, determine the initial volume of local anesthetic,  $Volume_{LA-initial}$ .
  - 10 ml
  - 11 ml
  - 12 ml
  - 13 ml
- Calculate the pump flow rate from Equations 6 or 7:
  - 0.4 ml/day
  - 0.5 ml/day
  - 0.6 ml/day
  - 0.7 ml/day
- Find the duration of the mixture in the reservoir from Equation 8.
  - 25 days
  - 30 days
  - 35 days
  - 40 days
- Use Equation 16 to determine that the initial and final amounts of local anesthetic, in milligrams, are equivalent.
  - 70 mg
  - 80 mg
  - 90 mg
  - 100 mg
- Use Equation 17 to determine that the initial and final amounts of morphine, in milligrams, are equivalent.
  - 130 mg
  - 140 mg
  - 150 mg
  - 160 mg

For a *fixed-rate* intrathecal pump, it is desired to deliver 3 mg per day of bupivacaine and 5 mg per day of morphine. The pump has a flow rate of 2 ml/day and a reservoir capacity of 50 ml. The initial concentration of local anesthetic,  $C_{LA-initial}$ , is 7.5 mg/ml and the initial concentration of morphine,  $C_{MSO_4-initial}$ , is 10 mg/ml.

- Find the final concentration of morphine,  $C_{MSO_4-final}$ , from Equation 9.
  - 2.5 mg/ml
  - 3.5 mg/ml
  - 4.5 mg/ml
  - 5.5 mg/ml
- Calculate the final concentration of local anesthetic,  $C_{LA-final}$ , from Equation 10.
  - 0.5 mg/ml
  - 1.5 mg/ml
  - 2.5 mg/ml
  - 3.5 mg/ml
- Using Equation 11, determine the initial volume of local anesthetic,  $Volume_{LA-initial}$ .
  - 7.0 ml
  - 8.0 ml
  - 9.0 ml
  - 10.0 ml
- Using Equation 12, find the initial volume of morphine,  $Volume_{MSO_4-initial}$ .
  - 10.5 mg
  - 11.5 mg
  - 12.5 mg
  - 13.5 mg
- Determine the volume of preservative-free normal saline,  $Volume_{NS}$ , to add to the combination

## Appendix: Continuing Medical Education Questions—Continued

- of Volume<sub>LA-initial</sub> and Volume<sub>M<sub>SO</sub>-initial</sub> so that the final volume is equal to that of the pump reservoir. Use Equation 13.
- 24.5 ml
  - 25.5 ml
  - 26.5 ml
  - 27.5 ml
15. Use Equation 16 to determine that the initial and final amounts of local anesthetic, in milligrams, are equivalent.
- 75 mg
  - 85 mg
  - 95 mg
  - 105 mg
16. Use Equation 17 to determine that the initial and final amounts of morphine, in milligrams, are equivalent.
- 110 mg
  - 115 mg
  - 120 mg
  - 125 mg

## Answers to CME Questions for Tutorial 24, Vol. 6, Issue 3

1. e
2. d
3. b
4. c
5. a
6. c
7. e
8. a

9. a
10. e
11. e
12. c
13. c
14. e
15. e